

Post Cretaceous/Paleogene boundary recovery of the deep marine ecosystem recorded by benthic foraminiferal morphogroups

Michael A. Kaminski, Syouma Hikmahtiar, and Claudia G. Cetean
Acta Palaeontologica Polonica 70 (4), 2025: 641-648 doi:10.4202/app.01227.2024

The uppermost 2 meters of the Maastrichtian Scaglia Rossa Fm. (Gubbio, Italy) record a notable change in the proportions of morphogroups of deep-water agglutinated foraminifera beginning about 1.4 m below the boundary, where we observe a gradual decrease in the abundance of the morphotype M4 (infauna) and a concurrent increase in the morphotype M1 (suspension feeders). This trend suggests a reduction in the flux of organic matter to the sea floor began approximately 114 k.y. before the boundary event. The abundance of M4 shows an abrupt decline from around 13% below the Cretaceous/Paleogene boundary, to a minimum of 7% in the beds directly above the boundary. This trend suggests a modest, but temporary reduction in organic matter flux to the sea floor following the Cretaceous/Paleogene boundary event, though not in the category of a “Strangelove Ocean”. Morphotype M1 shows a marked decrease across the boundary, with values decreasing from >70% to around 11%. The lower Paleocene displays strong fluctuations in the M4 morphogroup. In the same interval we observe a rise in Morphotype M2 (epifaunal detritivores) from values below 10% in the Maastrichtian to over 50% in the basal Paleocene, and a short-lived maximum in Morphotype M3, indicating a decline in particulate organic matter input from suspension and a concurrent increase in bacterial-derived organic matter in the early recovery phase of the Paleocene. The deep-marine ecosystem as witnessed by deep-water agglutinated foraminiferal morphogroups shows a prolonged recovery, with M1 abundance returning to Maastrichtian values approximately 2.8 meters above the Cretaceous/Paleogene boundary clay, corresponding to an age of 810 k.y. after the event. These findings support previous research from calcareous nannofossil studies, which suggested a prolonged recovery of the marine food web following the Cretaceous/Paleogene boundary event.

Key words: Foraminifera, ecology, Strangelove Ocean, Cretaceous, Paleogene, Apennines, Italy.

Michael A. Kaminski [kaminski@kfupm.edu.sa; ORCID: <https://orcid.org/0000-0002-7344-5874>], Geosciences Department, King Fahd University of Petroleum & Minerals, Dhahran 31261, Saudi Arabia. Syouma Hikmahtiar [syouma.santoso@kfupm.edu.sa; ORCID: <https://orcid.org/0000-0002-8481-1884>], Geosciences Department, King Fahd University of Petroleum & Minerals, Dhahran 31261, Saudi Arabia. Claudia G. Cetean [ceteanc@yahoo.com; ORCID: <https://orcid.org/0009-0004-9882-0257>], Micropress Europe, al. Mickiewicza 30, 30-059 Kraków, Poland.

This is an open-access article distributed under the terms of the Creative Commons Attribution License (for details please see creativecommons.org), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

 [Full text \(883.8 kB\)](#) |

 [Supplementary file \(136.2 kB\)](#)