

## Triassic Dasycladaceae from Central Balkan and Forebalkan, Bulgaria

Zbigniew Kotański and Georgy A. Catalov

*Acta Palaeontologica Polonica* 18 (2), 1973: 186-208

A very rich assemblage of Middle Anisian (Pelsonian) physoporellas is described. It belongs to the *Physoporella praealpina* group (*Ph. praealpina*, *Ph. dissita* and *Ph. minutula*) and to the group of *Ph. pauciforata* (*Ph. pauciforata* var. *pauciforata*, var. *sulcata* and var. *undulata*). A new variety *Ph. praealpina* var. *sulcata* is proposed. *Ph. praealpina* group characterises the Prealpes and Bransonian series in the Western Alps, as well as the High-Tatric series and Slovakian Karst in the Carpathians. Besides, *Gyroporella cf. ampleforata* and *G. vesiculifera* are described. This last species is an index Norian form and testifies to the close palaeogeographical connections between the Bulgarian, the Dinaric and the South Alpine seas in the Norian stage.

This is an open-access article distributed under the terms of the Creative Commons Attribution License (for details please see [creativecommons.org](https://creativecommons.org)), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

 [Full text \(1,759.7 kB\)](#)