

## Taxonomic re-assessment and phylogenetic relationships of Miocene homonymously spiral-horned antelopes

Dimitris S. Kostopoulos

*Acta Palaeontologica Polonica* 59 (1), 2014: 9-29 doi: <http://dx.doi.org/10.4202/app.2011.0013>

A morphological reappraisal of Middle Miocene and Early Pliocene antelopes with homonymously twisted horncores (i.e., with the left horncore twisted clockwise from the base up) and an additional re-description of some critical Greek specimens allows a revision of the current taxonomy at the species and genus level. A parsimony analysis recognizes *Hispanodorcas*, *Oioceros*, *Samodorcas*, *Samotragus*, *Paraoioceros*, and *Urmiatherium* as distinct lineages of the subtribe Oiocerina. The latter emerged during the Astaracian (ca. 14–11.5 Ma), and radiated in Eurasia during the mid-late Vallesian (ca. 10–9 Ma), resulting in a great variety of genera with unique combinations of morpho-ecological adaptations. The suprageneric relationships of Oiocerina are poorly understood, but their closest ancestors possibly lie within or close to the Middle Miocene “hypsodontines”, and inside Antilopinae.

**Key words:** Mammalia, Bovidae, Oiocerina, systematics, phylogeny, zoogeography, ecology, Miocene, Greece.

Dimitris S. Kostopoulos [[dkostop@geo.auth.gr](mailto:dkostop@geo.auth.gr)], University of Thessaloniki, Department of Geology, Laboratory of Paleontology, 54124 Thessaloniki, Greece.

This is an open-access article distributed under the terms of the Creative Commons Attribution License (for details please see [creativecommons.org](http://creativecommons.org)), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

