

## On microstructures of columnals of some Paleozoic crinoids

Edward Głuchowski

*Acta Palaeontologica Polonica* 27 (1-4), 1982: 77-83

Scanning electron microscope was used to study columnals of crinoids of *Pisocrinus* (?) *costatus* (Shevchenko), *Schyschcatocrinus* sp., *Cyclocaudiculus gracilis* Gluchowski, *Rhysocamax cristata* Moore et Jeffords and *Platycrinites minor* Gluchowski from Silurian, Devonian and Carboniferous deposits of Poland. The presence of microstructures, interpreted as stereom  $\alpha$  and stereom  $\beta$  have been confirmed and the occurrence of what is known as "large" microstructures on the lateral surface of the columnals of *Schyschcatocrinus* sp. have been stated. Three types within the "large" microstructures have been distinguished, the first of them marked by a close spacing of round, oval or elongated pores, the second with round, less closely spaced pores and the third - having irregular, labyrinthic furrows and depressions.

**Key words:** crinoids, microstructures, Paleozoic.

This is an open-access article distributed under the terms of the Creative Commons Attribution License (for details please see [creativecommons.org](https://creativecommons.org)), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

 [Full text \(1,021.1 kB\)](#)