

## The Jurassic juvenile ammonites of the Jagua Formation, Cuba

Cyprian Kulicki and Andrzej Wierzbowski

*Acta Palaeontologica Polonica* 28 (3-4), 1983: 369-384

Early ontogenetic stages of some Ammonitina (mostly Perisphinctidae) are described from the Oxfordian Jagua Formation of western Cuba. Forty six percent of the sample are specimens attaining the nepionic swelling with more than four septa developed. Juvenile jaw apparatus found for the first time in specimens with only 2,25 whorls, displays primitive elements corresponding to conchorhynch and rhyncholit of nautiloids, litoceratids and phylloceratids. The fauna and sediment features evidence a low water energy and low oxygenated conditions at the bottom during the sedimentation. The juvenile ammonites occur together with numerous adults which are represented by corresponding micro- and macroconchs: it indicates for the proximity of a breeding place.

**Key words:** ammonites, juvenile shells, juvenile jaw apparatus, paleoecology, Jagua Fm., Oxfordian, Cuba.

This is an open-access article distributed under the terms of the Creative Commons Attribution License (for details please see [creativecommons.org](http://creativecommons.org)), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

 [Full text \(6,730.1 kB\)](#)