

## Mid-Cretaceous microbiostratigraphy and foraminifers of the NE margins of the Świętokrzyskie (Holy Cross) Mts., Poland

Danuta Peryt

*Acta Palaeontologica Polonica* 28 (3-4), 1983: 417-466

Mid-Cretaceous foraminifers of the northeastern Mesozoic margin of the Holy Cross Mts. have been described. Planktic foraminifer standard zones - the *Rotalipora appenninnica*, *Rotalipora brotzeni*, *Rotalipora cushmani*, *Praeglobotruncana helvetica* - have been distinguished in the profile of Ożarów. A local benthic zonation, with the *Lingulogavelinella frankei*, *Gavelinella schloenbachi*, *Lingulogavelinella spinosa*, *Lingulogavelinella formosa*, *Gavelinella baltica* and *Lingulogavelinella globosa* zones, has been established in this profile. Due to the strong stratigraphic condensation and mixed fauna, no boundaries between the zones mentioned could be determined in the Rachów anticline.

**Key words:** Foraminiferida, Albian, Cenomanian, Turonian, stratigraphy, Poland.

This is an open-access article distributed under the terms of the Creative Commons Attribution License (for details please see [creativecommons.org](https://creativecommons.org)), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

 [Full text \(3,194.5 kB\)](#)