

Middle Triassic Scleractinia from the Cracow-Silesia region, Poland

Elżbieta Morycowa

Acta Palaeontologica Polonica 33 (2), 1988: 91-121

Scleractinian coral buildups from the Karchowice Beds (uppermost part of the Lower Muschelkalk) and from the Diplopora Dolomite (Middle Muschelkalk) of the Cracow-Silesia region were examined. Described corals comprise: three species known from literature-*Coelocenia* cf. *decipiens* (Laube), *Pamiroseris silesiaca* (Beyrich), *Eckastraea prisca* (Weissermel); four new species -*Volzeia szulci*, *Morycastraea opoliensis*, *Silesiastraea weissermeli*, *Chevalieria tenuiseptata*; and three determined as *Stylophylloopsis* sp., *Cyathocoenia* sp. and *Pamiroseris* sp. New genera, *Eckastraea* and *Sileriastraea*, are established. The most frequent component of coral communities appears *Pamiroseris silesiaca*.

Key words: Scleractinia, taxonomy, morphology, paleoecology, Middle Triassic, southern Poland.

This is an open-access article distributed under the terms of the Creative Commons Attribution License (for details please see creativecommons.org), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

 [Full text \(2,323.0 kB\)](#)