

Ontogenetic development of the thecal structures in caryophylliine scleractinian corals

Jarosław Stolarski

Acta Palaeontologica Polonica 40 (1), 1995: 19-44

At the initial stage of ontogeny, in Caryophylliidae (Miocene *Canryophyllia salinario*, *C. depauperata*, Recent *C. berteriana*) and Flabellidae (Miocene *Flabellum roissyanum*, Recent *Javania cailleti*), wall and septa are formed simultaneously, and their trabecular structure is coalesced (marginothecal wall). At subsequent juvenile stage in Caryophylliidae the presence of the extensive exosarc enables formation of costo-septa and, in consequence, formation of trabeculotheca. Trabeculotheca consists of fragments of primordial wall located between the costosepta.

The trabeculothecal segments vanish in the adult stage in the majority of corals when the septothecal wall is formed by thickening of the costo-septa. In others, however, marginotheca can be present throughout the whole ontogenetic sequence (*C. salinaria*). Most Flabellidae are characterized by limited expression of exosarc and the presence of marginothecal wall up to the adult stage. The origin of 'flabellid' organization in Caryophylliina may result from a simple modification of ontogeny - extension of initial morphology to later ontogenetic stages. Such corals could develop several times, and the Flabellidae may be polyphyletic.

Key words: Scleractinia, wall structure, ontogeny, phylogeny

This is an open-access article distributed under the terms of the Creative Commons Attribution License (for details please see creativecommons.org), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.