

Retiolitid graptolite *Spinograptus* from Poland and its membrane structures

Anna Kozłowska-Dawidziuk

Acta Palaeontologica Polonica 42 (3), 1997: 391-412

Spinograptus reticulolawsoni sp. n., *S. latespinosus* sp. n., *S. cf. latespinosus*, and *S. sp.* are described from borecore samples of the late Wenlock (*Colonograptus? praedeubeli* Zone), of the Bartoszyce borehole, NE Poland. They represent an early diversification phase of retiolitids, which followed the *Cyrtograptus lundgreni* Event. Some species display a combination of primitive and advanced characters, implying a mosaic pattern of *Spinograptus* evolution. Most of the material studied contains continuous peridermal membranes, suggesting that they may have been universally present in the retiolitids. Long (2 mm) siculae found in two species of *Spinograptus* contrast with short siculae, known for *Retiolites* and *Paraplectograptus*.

Key words: graptolites, retiolitids, membrane, phylogeny, Silurian, Poland.

Anna Kozłowska-Dawidziuk [akd@twarda.pan.pl], Instytut Paleobiologii PAN, ul. Twarda51/55, PL-00-818 Warszawa, Poland.

This is an open-access article distributed under the terms of the Creative Commons Attribution License (for details please see creativecommons.org), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.