

## Latest Frasnian Atrypida (Brachiopoda) from South China

Ma Xueping

*Acta Palaeontologica Polonica* 43 (2), 1998: 345-360

In South China latest Frasnian (*Palmatolepis linguiformis* Zone) the representatives of the order Atrypida (Brachiopoda) are most common in central Hunan Province. They are relatively rare in other parts of South China due to unfavourable ecologic conditions. Unlike most previously reported sections, including some sections in South China, the four Frasnian-Famennian (F-D) boundary sections examined here do not show any evidence for black shale. Atrypids are abundant and relatively diverse about 15 to 20 m below the F-F boundary (six species), and very rare about 1-2 m below the boundary (with only two species). It seems that the disappearance of most atrypids occurred well before the F-F boundary. Nine species (including *lowatrypa? qidongensis* sp. n.), assigned to six genera, are discussed and described.

**Key words:** Brachiopoda, Atrypida, taxonomy, biostratigraphy, biogeography, mass -extinction, Frasnian, Devonian, South China.

Ma Xueping [[xma@geoms.geo.pku.edu.cn](mailto:xma@geoms.geo.pku.edu.cn)], Department of Geology, Peking University, Beijing 100871, China.

This is an open-access article distributed under the terms of the Creative Commons Attribution License (for details please see [creativecommons.org](http://creativecommons.org)), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.