

## The Late Devonian trematid lingulate brachiopod *Schizobolus* from Poland

Andrzej Baliński and Lars E. Holmer

*Acta Palaeontologica Polonica* 44 (3), 1999: 335-346

A new species of the poorly known lingulate brachiopod *Schizobolus* is described from the Famennian (Upper Devonian) of Poland. *S. polonicus* sp. n. has a triangular pedicle notch and a small listrium, indicating that it belongs to the Trematidae within the superfamily Discinoidea. *S. polonicus* retains some linguloid features, such as a linguloid-like 'pedicle groove' and a V-shaped imprint of the pedicle nerve. The disturbance band, which occurs in the apical part of the larval shell, probably delimits two stages of growth, namely pre-larval (embryonic?) and larval, or, early-larval and late-larval. *S. polonicus* is the youngest member of the genus, and of the family Trematidae. Five incompletely preserved discinids from the Famennian of Łagów are described as Trematidae gen. et sp. indet.

**Key words:** Brachiopoda, Discinoidea, systematics, larval shell, Late Devonian, Poland.

Andrzej Baliński [[balinski@twarda.pan.pl](mailto:balinski@twarda.pan.pl)], Instytut Paleobiologii PAN, ul.

Twarda 51/55, PL-00-818 Warszawa, Poland. Lars E. Holmer [[Lars.Holmer@pal.uu.se](mailto:Lars.Holmer@pal.uu.se)], Institute of Earth Sciences, Department of Historical Geology & Palaeontology, Norbyvägen 22, S-752 36 Uppsala, Sweden.

This is an open-access article distributed under the terms of the Creative Commons Attribution License (for details please see [creativecommons.org](http://creativecommons.org)), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.