

## Body mass estimation and locomotion of the Miocene pelecaniform bird *Macranhinga*

Jorge Ignacio Noriega

*Acta Palaeontologica Polonica* 46 (2), 2001: 247-260

The fossil darter *Macranhinga paranensis* (Aves: Pelecaniformes) from the late Miocene of Argentina is the largest of all known extinct or living Anhingidae. Its body mass is estimated at a mean value of 5.4 kg by using a scaling model derived from the logarithmic relationship between measurements of the least shaft circumference of the femur/ tibiotarsus and body masses. Predictions of body mass, as well as the analysis of anatomical evidence, are used to infer that *Macranhinga paranensis* would have probably had a powered flapping flight and an aquatic locomotion similar to that of cormorants. The morphology of the pelvis and the hind-limb would have allowed *Macranhinga paranensis* to catch fishes by means of pursuit-diving rather than stalking them in an anhinga-like manner. As determined by adaptations mainly of tarsometatarsal morphology, the species had well developed perching and climbing abilities.

**Key words:** Aves, Anhingidae, *Macranhinga*, Paleobiology, Miocene, Argentina.

Jorge Ignacio Noriega [[cidnoriega@infoshopde.com.ar](mailto:cidnoriega@infoshopde.com.ar)], Centro de Investigaciones Científicas y de Transferencia Tecnológica - CONICET. Dr. Materi y España, (3105) Diamante, provincia de Entre Ríos, Argentina.

This is an open-access article distributed under the terms of the Creative Commons Attribution License (for details please see [creativecommons.org](http://creativecommons.org)), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.