

## Aalenian to Cenomanian terrestrial palynofloras of SW Scania, Sweden

Vivi Vajda

*Acta Palaeontologica Polonica* 46 (3), 2001: 403-426

I describe dispersed miospore assemblages recovered from 35 drill-core samples from Höllviken 2, Norrevíng 1, and Svedala 1 wells, all in SW Scania, Sweden. Over eighty taxa of pollen and spores, ranging from the Aalenian to the Cenomanian were identified. Four pollen/spore zones have been defined on the basis of key taxa and on the variation in the frequency of miospore groups. The palynofacies analysis indicates that a continental depositional environment prevailed during the Aalenian in Scania. A stratigraphic hiatus existed from the Aalenian then on until the Valanginian, when the depositional environment subsequently became marine. The marine conditions continued until the Cenomanian. A gradual increase in marine palynomorphs is found in the Cenomanian succession, indicating a transgression. On the basis of the palynoflora it is suggested that the vegetation consisted of cycads, conifers, pteridophytes and a very limited number of angiosperms.

**Key words:** Palynology, paleovegetation, paleoclimate, Cretaceous, sealevel-changes, Scania, Sweden.

Vivi Vajda [[vivi.vajda@geol.lu.se](mailto:vivi.vajda@geol.lu.se)], Department of Geology, Division of Historical Geology and Palaeontology, University of Lund, Sölvegatan 13, SE-223 62 Lund, Sweden.

This is an open-access article distributed under the terms of the Creative Commons Attribution License (for details please see [creativecommons.org](http://creativecommons.org)), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.