

## A new docodont mammal from the Late Jurassic of the Junggar Basin in Northwest China

Hans-Ulrich Pfretzschner, Thomas Martin, Michael W. Maisch, Andreas T. Matzke, and Ge Sun  
*Acta Palaeontologica Polonica* 50 (4), 2005: 799-808

Fieldwork in the early Late Jurassic (Oxfordian) Qigu Formation of the Junggar Basin in Northwest China (Xinjiang Autonomous Region) produced teeth and mandibular fragments of a new docodont. The new taxon has a large 'pseudotalonid' on the lower molars, and by retention of crest b-g exhibits closer affinities to *Simpsonodon* and *Krusatodon* from the Middle Jurassic of Europe than to the other known Asian docodonts *Tashkumyrodon*, *Tegotherium*, and *Sibirotherium*. It differs from the *Haldanodon-Docodon*-lineage by the 'pseudotalonid' and large cusps b and g. A PAUP analysis based on lower molar characters produced a single most parsimonious tree with two main clades. One clade comprises *Docodon*, *Haldanodon*, and *Borealestes*, and the other *Dsungarodon*, *Simpsonodon*, and *Krusatodon* plus the Asian tegotheriids. Analysis of the molar occlusal relationships using epoxy casts mounted on a micromanipulator revealed a four-phase chewing cycle with transverse component. The molars of the new docodont exhibit a well developed grinding function besides cutting and shearing, probably indicating an omnivorous or even herbivorous diet. A grinding and crushing function is also present in the molars of *Simpsonodon*, *Krusatodon*, and the Asian tegotheriids, whereas *Borealestes*, *Haldanodon*, and *Docodon* retain the plesiomorphic molar pattern with mainly piercing and cutting function.

**Key words:** Docodonta, *Dsungarodon*, occlusion, Jurassic, Qigu Formation, Junggar Basin.

Hans-Ulrich Pfretzschner [[hans-ulrich.pfretzschner@uni-tuebingen.de](mailto:hans-ulrich.pfretzschner@uni-tuebingen.de)] and Michael W. Maisch [[michael.maisch@uni-tuebingen.de](mailto:michael.maisch@uni-tuebingen.de)], Institut für Geowissenschaften, Universität Tübingen, Sigwartstraße 10, 72076 Tübingen, Germany; Thomas Martin [[tmartin@senckenberg.de](mailto:tmartin@senckenberg.de)], Forschungsinstitut Senckenberg, Senckenberganlage 25, 60325 Frankfurt am Main, Germany; Andreas T. Matzke [[matzkea@si.edu](mailto:matzkea@si.edu)], Department of Paleobiology, Smithsonian Institution, National Museum of Natural History, PO Box 37012 MRC 121, Washington, D.C. 20013-7012, USA; Ge Sun [[sunge@jlu.edu.cn](mailto:sunge@jlu.edu.cn)], Research Center of Paleontology, Jilin University, 6, Xi-Minzhū Street, Changchun 130026, China.

This is an open-access article distributed under the terms of the Creative Commons Attribution License (for details please see [creativecommons.org](https://creativecommons.org)), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

 [Full text \(428.2 kB\)](#)