

A new heterosoricid shrew from the lowermost Oligocene of Europe

Richard Smith and Lars W. van den Hoek Ostende
Acta Palaeontologica Polonica 51 (2), 2006: 381-384

The earliest Oligocene faunas of Europe are characterised by a large number of Asian immigrants. One of the classical sites that shows the faunal change after this so-called “Grande Coupure” is Hoogbutsel in Belgium (MP 21). Recently a new locality from the lowermost Oligocene was discovered in Belgium, Boutersem TGV. Here, we describe a new heterosoricid (Mammalia, Soricomorpha), *Belgicasorex ramboeri* gen. et sp. nov., from these lowermost Oligocene localities. We assume that, like the erinaceid *Tetracus nanus* Aymard, 1846 and the nyctitheriid *Oligonyctia hoffmani* Smith, 2004, *Belgicasorex ramboeri* was one of the Asian immigrants that entered Europe after the “Grande Coupure”.

Richard Smith [richardsmithpal@hotmail.com], Département de Paléontologie, Institut royal des Sciences naturelles de Belgique, rue Vautier 29, B-1000 Bruxelles, Belgium; Lars W. van den Hoek Ostende [Hoek@Naturalis.nl], Nationaal Natuurhistorisch Museum, Naturalis, Postbus 9517, NL-2300 RA Leiden, The Netherlands.

This is an open-access article distributed under the terms of the Creative Commons Attribution License (for details please see creativecommons.org), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.