

## The Barremian heteromorph ammonite *Dissimilites* from northern Italy: Taxonomy and implications

Alexander Lukeneder and Susanne Lukeneder

*Acta Palaeontologica Polonica* 59 (3), 2014: 663-680 doi: <http://dx.doi.org/10.4202/app.2012.0014>

A new acrioceratid ammonite, *Dissimilites intermedius* sp. nov., from the Barremian (Lower Cretaceous) of the Puez area (Dolomites, northern Italy) is described. *Dissimilites intermedius* sp. nov. is an intermediate form between *D. dissimilis* and *D. trinodosum*. The new species combines the ribbing style of *D. dissimilis* (bifurcating with intercalating single ribs) with the tuberculation style of *D. trinodosum* (trituberculation on entire shell). The shallow-helical spire, entirely comprising single ribs intercalated by trituberculated main ribs, is similar to the one of the assumed ancestor *Acrioceras*, whereas the increasing curvation of the younger forms resembles similar patterns observed in the descendant *Toxoceratoides*. These characters support the hypothesis of a direct evolutionary lineage from *Acrioceras* via *Dissimilites* to *Toxoceratoides*. *D. intermedius* sp. nov. ranges from the upper Lower Barremian (*Moutoniceras moutonianum* Zone) to the lower Upper Barremian (*Toxancyloceras vandenheckii* Zone). The new species allows to better understand the evolution of the genus *Dissimilites*. The genus appears within the *Nicklesia pulchella* Zone represented by *D. duboise*, which most likely evolved into *D. dissimilis*. In the *Kotetishvilia compressissima* Zone, two morphological forms developed: smaller forms very similar to *Acrioceras* and forms with very long shaft and juvenile spire like in *D. intermedius* sp. nov. The latter most likely gave rise to *D. subalternatus* and *D. trinodosum* in the *M. moutonianum* Zone, forms which were probably ancestral to the genus *Toxoceratoides*.

**Key words:** Ammonoidea, Acrioceratidae, *Dissimilites*, Barremian, Cretaceous, Alps, Italy.

Alexander Lukeneder [[alexander.lukeneder@nhm-wien.ac.at](mailto:alexander.lukeneder@nhm-wien.ac.at)], Geological-Palaeontological Department, Natural History Museum, Burgring 7, 1010 Vienna, Austria; Susanne Lukeneder [[susanne.lukeneder@nhm-wien.ac.at](mailto:susanne.lukeneder@nhm-wien.ac.at)], Geological-Palaeontological Department, Natural History Museum, Burgring 7, 1010 Vienna and Department of Palaeontology, University of Vienna, Althanstrasse 14, 1090 Vienna, Austria.

This is an open-access article distributed under the terms of the Creative Commons Attribution License (for details please see [creativecommons.org](https://creativecommons.org)), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

 [Full text \(842.8 kB\)](#)