

## **Givetian (Middle Devonian) sharks from Cairo, New York (USA): Evidence of early cosmopolitanism**

Daniel Potvin-Leduc, Richard Cloutier, Ed Landing, Linda VanAller Hernick, and Frank Mannolini *Acta Palaeontologica Polonica* 60 (1), 2015: 183-200 doi: http://dx.doi.org/10.4202/app.2012.0101

Whereas cosmopolitan distribution patterns are established for many Late Devonian vertebrates (e.g., placoderms, onychodontiforms), few palaeobiogeographic studies have considered chondrichthyans. Recent discoveries of shark material demonstrate that some chondrichthyans were cosmopolitan by the Middle Devonian. Abundant Givetian microremains have been recovered from the Cairo quarry in eastern New York State, USA. These include teeth of two shark species with Gondwanan affinities, the omalodontid Portalodus mannoliniae sp. nov. and the antarctilamnid Wellerodus priscus. Abundant teeth of P. mannoliniae sp. nov. are characterized by a smooth diplodont crown, polarized cusps, and a labially oriented base. The teeth demonstrate monognathic heterodonty. The juvenile morph is distinguished from the adult by smaller size, slender cusps, and variation in the shape of the base. W. priscus is represented by rare juvenile teeth. Two groups of scales that show affinity to material from northern (Spain) and East Gondwana (Antarctica) are tentatively attributed to the two described species. Antarctilamnid distribution suggests a north Gondwanan origin and a colonization of the margin of the landmass before dispersing to Laurentia by the Middle Devonian. This material further indicates that vertebrate global dispersal was initiated by the Middle Devonian, and emphasizes earlier palaeogeographic interpretations that the Middle Devonian "Hamilton fauna" of North American Laurussia originated in the Early Devonian in South American Gondwana.

**Key words:** Chondrichthyes, Elasmobranchii, Antarctilamnidae, Omalodontiformes, taxonomy, ontogeny, palaeobiogeography, Devonian, North America.

Daniel Potvin-Leduc [potvinleduc@gmail.com] and Richard Cloutier [richard\_cloutier@uqar.ca ], Laboratoire de Biologie évolutive, Université du Québec à Rimouski, 300 allée des Ursulines, Rimouski, Québec, G5L 3A1, Canada; Ed Landing [elanding@mail.nysed.gov ], Linda VanAller Hernick [lhernick@mail.nysed.gov], and Frank Mannolini [fmannoli@mail.nysed.gov], New York State Museum, 222 Madison Avenue, Albany, New York 12230, USA. This is an open-access article distributed under the terms of the Creative Commons Attribution License (for details please see <u>creativecommons.org</u>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

