

Turonian and Coniacian foraminifera from the Nysa Trough, Sudetes, Poland

Eugenia Gawor-Biedowa

Acta Palaeontologica Polonica 25 (1), 1980: 3-54

Turonian to Coniacian foraminifera from the Nysa Trough, Sudetes, are investigated. Totally, 55 species are recorded; 42 species in the Turonian, 49 species in the Coniacian. Most species are benthic forms. Two new species are described from the Turonian: *Dicarinella sudetica* sp. n. and *Dicarinella radwanskae* sp. n. Wall microstructure and alternation of generations are studied in the genus *Gavetina*. Six foraminiferal zones are recognized; 4 of them in the Turonian (*Dicarinella sudetica* Zone, *Tappanina eouvigeriniformis* Zone, *Archaeoglobigerina cretacea* Zone, and *Dicarinella concavata* Zone), and the remaining 2 in the Coniacian (*Epistomina spinulifera potyptioides* Zone and *Gaudryina sudetica* Zone). Changes in composition of the foraminiferal assemblages were caused by changes in water depth and presence/absence of interconnections between the basin and open sea.

Key words: Foraminifera, Turonian, Coniacian, stratigraphy, Sudetes, Poland.

This is an open-access article distributed under the terms of the Creative Commons Attribution License (for details please see creativecommons.org), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.