

Orthid brachiopods from the Upper Visean (Carboniferous) of the Świętokrzyskie Mts., Poland

Halina Żakowa

Acta Palaeontologica Polonica 34 (2), 1989: 91-124

Enteletid (four species) and rhipidomellid (one species) brachiopoda from the Gałęzice region, SW Świętokrzyskie Mts., have been described. Dominating genus *Schizophoria* is represented by *S. (Schizophoria) resupinata*, *S. (S.) keyserlingiana*, *S. (Paraschizophoria) woodi*, *S. (Pat.) sp. A*, *S. (Pocockia) linguata*, *S. (P.) cf. gibbera*. Six known morphological "varieties" or "forms" of *S. (S.) resupinata* have been described. They probably come from diverse environments. Most specimens collected belong to *S. (S.) resupinata* (Martin) forma typica and represent all stages of its ontogeny. The brachiopods have been collected from organodetritic limestones and calcareous-clayey deposits of the *Goniatites crenistria* and *G. striatus* zones. It is suggested that only *S. (S.) resupinata* f. *gigantea* may be stratigraphically characteristic of the Upper Visean of Europe.

Key words: brachiopods, taxonomy, stratigraphy, Upper Visean, Świętokrzyskie Mts., Poland.

This is an open-access article distributed under the terms of the Creative Commons Attribution License (for details please see creativecommons.org), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.