

Brachiopods and conodonts from the Early Carboniferous of South China

Andrzej Baliński

Acta Palaeontologica Polonica 44 (4), 1999: 437-451

A sample taken from a detrital limestone lens, presumed to be allochthonous, within the dark coloured argillaceous limestone of the Early Carboniferous Muhua Formation at the Muhua section, Guizhou, South China, yielded numerous, mostly silicified fossils. Ostracodes, which are the most numerous in the sample, were studied by Olempska (1999). Brachiopods and conodonts are described and illustrated in this paper, but other associated fossils are also noted. Among brachiopods the most common are productides, orthotetidines, spiriferides, and orthides. The productoid gen. et sp. indet. 2, *Lambdarina* sp., and rhynchonelloid gen. et sp. indet. most probably represent new taxa, but are described in open nomenclature because of inadequate material. Conodonts are indicative of late Tournaisian age. The fossil assemblage is represented by phosphatic and silicified remnants, the latter being originally calcitic. The pattern of silicification resulted generally in preservation of skeletal morphology in great details.

Key words: Brachiopoda, Conodonta, Early Carboniferous, silicification, China.

Andrzej Baliński [balinski@twarda.pan.pl], Instytut Paleobiologii PAN, ul. Twarda 51/55, PL-00-818 Warszawa, Poland.

This is an open-access article distributed under the terms of the Creative Commons Attribution License (for details please see creativecommons.org), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.