

## New data on the Paleocene monotreme *Monotrematum sudamericanum*, and the convergent evolution of triangulate molars

Rosendo Pascual, Francisco J. Goin, Lucía Balarino, and Daniel E. Udrizar Sauthier  
*Acta Palaeontologica Polonica* 47 (3), 2002: 487-492

We describe an additional fragmentary upper molar and the first lower molar known of *Monotrematum sudamericanum*, the oldest Cenozoic (Paleocene) monotreme. Comparisons suggest that the monotreme evolution passed through a stage in which their molars were "pseudo-triangulate", without a true trigonid, and that the monotreme pseudo-triangulate pattern did not arise through rotation of the primary molar cusps. Monotreme lower molars lack a talonid, and consequently there is no basin with facets produced by the wearing action of a "protocone"; a cristid obliqua connecting the "talonid" to the "trigonid" is also absent. We hypothesize that acquisition of the molar pattern seen in *Steropodon galmani* (Early Cretaceous, Albian) followed a process similar to that already postulated for docodonts (*Docodon* in Laurasia, *Reigitherium* in the South American sector of Gondwana) and, probably, in the gondwanathere *Ferugliotherium*.

**Key words:** Monotremata, Monotrematum, pseudo-triangulate molars, molar structure, Gondwana, Patagonia, Paleocene

Rosendo Pascual [[ropascua@museo.fcnym.unlp.edu.ar](mailto:ropascua@museo.fcnym.unlp.edu.ar)], Francisco Goin [[fgoin@museo.fcnym.unlp.edu.ar](mailto:fgoin@museo.fcnym.unlp.edu.ar)], Lucía Balarino, and Daniel Udrizar Sauthier, Departamento Paleontología Vertebrados, Museo de la Plata, Paseo del Bosque s/n, 1900 La Plata, Argentina.

This is an open-access article distributed under the terms of the Creative Commons Attribution License (for details please see [creativecommons.org](http://creativecommons.org)), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

