

Crested antiarch *Bothriolepis zadonica* H.D. Obrucheva from the Lower Famennian of Central European Russia

Sergey Moloshnikov

Acta Palaeontologica Polonica 49 (1), 2004: 135-146

New materials on the bothriolepidid *Bothriolepis zadonica* H.D. Obrucheva, 1983 from the Zadonskian Regional Stage (Upper Devonian, Lower Famennian) of Central Devonian Field (Central European Russia) are described and illustrated. These materials came from nine localities in the Orel and Tver Regions of Russia. This species is characterised by the presence of a well developed median dorsal crest in the trunk armour and unusual type of the preorbital recess of the head shield; this recess is designated herein 'trapezoid'. The ontogenetic and individual variations of *B. zadonica* are studied. The crested bothriolepidids are known only from Euramerica and East Gondwana. The possible relationships between crested bothriolepidids in these provinces are discussed.

Key words: Placodermi, Bothriolepidoidei, *Bothriolepis*, Devonian, Famennian, Russia.

Sergey Moloshnikov [molsergey@rambler.ru], Department of Palaeontology, Geological Faculty, Lomonosov Moscow State University, Vorob'evye Gory, 119992, Moscow, Russia.

This is an open-access article distributed under the terms of the Creative Commons Attribution License (for details please see creativecommons.org), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.