

## A peculiar faunivorous metatherian from the early Eocene of Australia

Robin M.D. Beck

*Acta Palaeontologica Polonica* 60 (1), 2015: 123-129 doi: <http://dx.doi.org/10.4202/app.2013.0011>

I describe *Archaeonothos henkgodthelpi* gen. et. sp. nov., a small (estimated body mass ~40–80 g) tribosphenic metatherian from the early Eocene Tingamarra Fauna of southeastern Queensland, Australia. This taxon, known only from a single isolated upper molar (M2 or M3) is characterised by a very distinctive combination of dental features that, collectively, probably represent faunivorous adaptations. These include: a straight, elevated centrocrista; a metacone considerably taller than the paracone; a wide styler shelf (~50% of the total labiolingual width of the tooth); reduced styler cusps; a long postmetacrasta; a small and anteroposteriorly narrow protocone; an unbasined trigon; and the absence of conules. Some of these features are seen in dasyuromorphians, but detailed comparisons reveal key differences between *A. henkgodthelpi* and all known members of this clade. *A. henkgodthelpi* also predates recent molecular estimates for the divergence of crown-group Dasyuromorphia. Similar dental features are seen in a number of other metatherians, including the South American sparassodonts, *Wirunodon chanku* from the ?middle–late Eocene Santa Rosa local fauna of Peru, and *Kasserinotherium tunisiense* from the early Eocene Chambi fauna of Tunisia, although whether *A. henkgodthelpi* is closely related to any of these taxa is unclear based on available evidence. I therefore refer *A. henkgodthelpi* to Metatheria incertae sedis. Potential relatives of *A. henkgodthelpi* are unknown from any other Australian fossil deposit.

**Key words:** Mammalia, Metatheria, Marsupialia, Sparassodonta, Eocene, Tingamarra Fauna, Australia.

Robin M.D. Beck [[r.m.d.beck@salford.ac.uk](mailto:r.m.d.beck@salford.ac.uk)], School of Environment & Life Sciences, Room G48, Peel Building, University of Salford, Salford M5 4WT, UK; and School of Biological, Earth and Environmental Sciences, University of New South Wales, Sydney, NSW 2052, Australia.

This is an open-access article distributed under the terms of the Creative Commons Attribution License (for details please see [creativecommons.org](http://creativecommons.org)), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

 [Full text \(207.2 kB\)](#)