

Some terebratulid populations from the Lower Kimmeridgian of Poland and their relations to the biotic environment

Jerzy Dzik

Acta Palaeontologica Polonica 24 (4), 1979: 473-492

Populations of *Epithyris* '*subsella*' in different Lower Kimmeridgian fossil assemblages show different size-frequency distributions although they exhibit a similar shape of survivorship curves. A population with large-sized adult specimens of *E.* '*subsella*' was characteristic for a community in which most abundant were bivalves: *Isognomon subplana*, *Lopha gregarea* and *Trichites saussurei*. A population with small adult specimens was characteristic for a community with abundant dasyclad alga *Goniolina geometrica*. Here, among shelled suspension-feeders the most important producers were brachiopods *E.* '*subsella*', *Zeilleria humeralis*, and *Septaliphoria pinguis*. Thalli of *G. geometrica* were the substrate for most of the sedentary organisms in this community. The estimation of biomass and productivity of particular species in fossil communities is discussed.

Key words: Brachiopods, fossil assemblages, Upper Jurassic, Poland.

This is an open-access article distributed under the terms of the Creative Commons Attribution License (for details please see creativecommons.org), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

 [Full text \(1,085.4 kB\)](#)