

The significance of graptoblasts in the life cycle of crustoid graptolites

Adam Urbanek

Acta Palaeontologica Polonica 28 (3-4), 1983: 313-326

Graptoblasts are small ovoid bodies with a flat lower, and a convex upper, wall, the latter provided with a costulation having a fusellar aspect. First found as isolated bodies associated with graptolite remains, graptoblasts were later

recognized as a constituent of crustoid colonies (Kozłowski 1949, 1962). Their biological role remain largely enigmatic. The view that graptoblasts were formed within the authothecae is rejected and a conclusion is advanced that

graptoblasts were closed, resting terminal portions of the stolothecae, housing eneysted dormant zooids. They may be compared with the resting terminal zooids in *Rhabdopleura* and with the hibernacula of ctenostome bryozoans. Graptoblasts provided an adaptation allowing the species to survive the periods of adverse conditions when the rest of the colony disintegrated. One could hypothesize that after germination the graptoblasts produced small propagules ejected through a narrow cryptopyle and forming new colonies after they settled on the substrate.

Key words: Graptoblasts, Graptolithina, Crustoidea, dormancy, germination, propagules, life cycle.

This is an open-access article distributed under the terms of the Creative Commons Attribution License (for details please see creativecommons.org), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.



[Full text \(849.8 kB\)](#)