

## A new species of *Glyphea* (Decapoda: Palinura) from the La Meseta Formation (Eocene) of Seymour Island, Antarctica

Rodney M. Feldmann and Andrzej Gaździcki

*Acta Palaeontologica Polonica* 42 (3), 1997: 437-445

A new species of palinuran lobster, *Glyphea reticulata*, from the lowermost part of the Eocene La Meseta Formation on Seymour Island, Antarctica, represents one of the stratigraphically youngest species of *Glyphea*. The occurrence of the last vestiges of what was previously a cosmopolitan genus in a region dominated by Pacific Ocean faunal influences is significant because the sole extant species of the Glypheidae, *Neoglyphea inopinata* Forest & Saint Laurent, 1975, is known only from the west Pacific.

**Key words:** Decapoda, Glypheidae, paleobiogeography, evolution, Eocene, Antarctica.

Rodney M. Feldmann [[rfeldman@kentvm.kent.edu](mailto:rfeldman@kentvm.kent.edu)], Department of Geology, Kent State University, Kent, Ohio, 44242, U.S.A. Andrzej Gaździcki [[gazdzick@twarda.pan.pl](mailto:gazdzick@twarda.pan.pl)], Instytut Paleobiologii PAN, ul. Twarda 51/55, PL-00-818 Warszawa, Poland.

This is an open-access article distributed under the terms of the Creative Commons Attribution License (for details please see [creativecommons.org](http://creativecommons.org)), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.