

Brachiopods from the Middle Ordovician Shihtzupu Formation of Yunnan Province, China

Renbin Zhan and Jisuo Jin Acta Palaeontologica Polonica 50 (2), 2005: 365-393

The brachiopod fauna of the Middle Ordovician (upper Darriwilian) Shihtzupu Formation in the Weixin area, northeastern Yunnan Province, southwestern China, comprises 16 genera and 18 species, including one new genus (*Halirhachis*) and five new species (*Glyptorthis sarcina* , *Protoskenidioides weixinensis*, *Halirhachis leonina*, *Leptellina spatiosa*, and *Leptestiina veturna*). Cluster and principal component analyses of the latest Arenigian-mid Caradocian faunas of the Upper Yangtze Platform with selected faunas of similar age from other palaeoplates or terranes indicate that the Weixin brachiopods have closest faunal affinities to those of the typical Shihtzupu Formation in Guizhou Province and the Naungkangyi Group of Burma (Myanmar). During latest Arenigian-mid Caradocian times, the brachiopods of the Upper Yangtze Platform, Sibumasu, and Chu-Ili palaeogeographical regions constituted a distinct faunal province, characterized by a large number of endemic taxa as well as regionally widespread genera such as *Saucrorthis*, *Martellia*, and *Yangtzeella*. This brachiopod faunal province has very low similarity coefficients with the coeval brachiopod faunas of Laurentia and Avalonia.

Key words: Brachiopoda, taxonomy, palaeobiogeography, Ordovician, Darriwilian, southwestern China.

Renbin Zhan [<u>rbzhan@nigpas.ac.cn</u>], Nanjing Institute of Geology and Palaeontology, Chinese Academy of Sciences, Nanjing 210008, China; Jisuo Jin [<u>jjin@uwo.ca</u>], Department of Earth Sciences, University of WesternOntario, London, OntarioN6A 5B7, Canada.

This is an open-access article distributed under the terms of the Creative Commons Attribution License (for details please see <u>creativecommons.org</u>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

