

## Crouching theropod and *Navahopus* sauropodomorph tracks from the Early Jurassic Navajo Sandstone of USA

Jesper Milàn, David B. Loope, and Richard G. Bromley

*Acta Palaeontologica Polonica* 53 (2), 2008: 197-205 doi:<http://dx.doi.org/10.4202/app.2008.0203>

Numerous tracks and trackways are preserved in the a cross-strata of the Lower Jurassic Navajo Sandstone of northern Arizona and southern Utah, USA. Tracks and trackways of small theropod dinosaurs are particularly abundant within one 10-m-thick interval. This paper describes a crouching trace from a theropod dinosaur that shows impressions of all four limbs, the ischial callosity, the tail, and tracks leading to and away from the crouching site, and revises the interpretation of a well preserved trackway hitherto referred to the synapsid ichnogenus *Brasilichnium* and here considered to be from a sauropodomorph dinosaur. It is named *Navahopus coyoteensis* isp. nov. on the basis of morphological differences from the type ichnospecies *N. falcipollex*. The ichnofamily Navahopodidae is revised to include *Tetrasauropous unguiferus*, *Navahopus falcipollex*, and *N. coyoteensis*.

**Key words:** *Navahopus*, Navahopodidae, Sauropodomorpha, Theropoda, ichnology, locomotory habits, crouching trace

Jesper Milàn [[Milan@geol.ku.dk](mailto:Milan@geol.ku.dk)] and Richard G. Bromley [[rullard@geol.ku.dk](mailto:rullard@geol.ku.dk)],  
Department of Geography and Geology, University of Copenhagen, Øster Voldgade  
10, DK-1350 Copenhagen K, Denmark; David B. Loope [[dloope1@unl.edu](mailto:dloope1@unl.edu)], Department of  
Geosciences, University of Nebraska, Lincoln, NE 68588-0340, Lincoln, Nebraska, USA.

This is an open-access article distributed under the terms of the Creative Commons Attribution License (for details please see [creativecommons.org](http://creativecommons.org)), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

